

“Black hole Quasinormal modes”

Miok Park

Center for Theoretical Physics of Universe (CTPU-PTC),
Institute for Basic Science (IBS)

"2025 Summer school for Numerical Relativity and Gravitational waves"
@ KASI, Daejeon, S. Korea

July 30, 2025

Content

1. The aim of this lecture

normal modes vs quasi-normal modes

2. Quantum mechanics : A particle in a box

Schrödinger equation

A particle in a box : eigenvalues and eigenfunctions

Summary

3. Quantum field theory on curved spacetime : A particle in a black hole

Klein-Gordon equation

a particle in a black hole : eigenvalues and eigenfunctions

WKB method

Green's function method

Summary

4. Gravitational waves and QNM

Black Hole Ringdown

normal modes vs quasi-normal modes

The purpose of this lecture is to understand

normal modes vs **quasi-normal** modes

In this lecture, we will understand them by studying


- ▶ **normal** modes : a particle in a box
- ▶ **quasi-normal** modes : a particle in a black hole

quasi-

1. cf. apparently but not really, seemingly
2. cf. being partly or almost


normal modes vs quasi-normal modes

A particle in a box




normal modes vs quasi-normal modes

A particle in a box




normal modes vs quasi-normal modes

A particle in a box




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box




A particle in a black hole




normal modes vs quasi-normal modes

A particle in a box

Closed system, energy is conserved!




A particle in a black hole



normal modes vs quasi-normal modes


A particle in a box

Closed system, energy is conserved!



A particle in a black hole

Open system, energy is NOT conserved!



Quantum mechanics

Schrödinger equation

- ▶ the time-dependent Schrödinger equation

$$i\hbar \frac{\partial \psi(x, t)}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi(x, t)}{\partial x^2} + V(x)\psi(x, t) \equiv \hat{H}\psi(x, t),$$

where $\psi(x, t)$: wave function

- ▶ If V has no explicit time dependence, we can make separation of variables

$$\psi(x, t) = T(t)u(x),$$

$$i\hbar \frac{1}{T(t)} \frac{dT(t)}{dt} = \frac{1}{u(x)} \left\{ -\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) \right\},$$

$$i\hbar \frac{dT(t)}{dt} = ET(t), \quad T(t) = Ce^{-iEt/\hbar},$$

- ▶ the time-independent Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = \hat{H}u(x) = Eu(x)$$

Hermitian (self-adjoint) operator

- ▶ In QM, **observables** correspond to **operators** that have certain special properties.

Hermitian (self-adjoint) operator

- ▶ In QM, **observables** correspond to **operators** that have certain special properties.
- A. the expectation value of an **observable** must be a **real number**

$$\langle \hat{O} \rangle = \int \psi^* \hat{O} \psi dx = \langle \psi | \hat{O} \psi \rangle = \langle \hat{O} \rangle^* \quad (1)$$

Hermitian (self-adjoint) operator

- In QM, **observables** correspond to **operators** that have certain special properties.
- A. the expectation value of an **observable** must be a **real number**

$$\langle \hat{O} \rangle = \int \psi^* \hat{O} \psi dx = \langle \psi | \hat{O} \psi \rangle = \langle \hat{O} \rangle^* \quad (1)$$

- B. for **operators** that represent **observables**

$$\langle \psi, \hat{O} \psi \rangle = \langle \hat{O} \psi, \psi \rangle \quad \left(\langle g, f \rangle = \langle f, g \rangle^*, \psi(x) \in \mathcal{H} \right) \quad (2)$$

$$\text{Proof :} \quad \hat{O} \psi = E \psi, \quad (3)$$

$$\langle f, \hat{O} f \rangle = \langle f, \hat{O} f \rangle^* = \langle \hat{O} f, f \rangle, \quad (4)$$

$$E \langle f, f \rangle = E^* \langle f, f \rangle \quad (5)$$

where $\langle f, f \rangle$ cannot be zero, so $E = E^*$ and hence E is real.

Hermitian (self-adjoint) operator

- In QM, **observables** correspond to **operators** that have certain special properties.

A. the expectation value of an **observable** must be a **real number**

$$\langle \hat{O} \rangle = \int \psi^* \hat{O} \psi dx = \langle \psi | \hat{O} \psi \rangle = \langle \hat{O} \rangle^* \quad (1)$$

B. for **operators** that represent **observables**

$$\langle \psi, \hat{O} \psi \rangle = \langle \hat{O} \psi, \psi \rangle \quad \left(\langle g, f \rangle = \langle f, g \rangle^*, \psi(x) \in \mathcal{H} \right) \quad (2)$$

$$\text{Proof :} \quad \hat{O} \psi = E \psi, \quad (3)$$

$$\langle f, \hat{O} f \rangle = \langle f, \hat{O} f \rangle^* = \langle \hat{O} f, f \rangle, \quad (4)$$

$$E \langle f, f \rangle = E^* \langle f, f \rangle \quad (5)$$

where $\langle f, f \rangle$ cannot be zero, so $E = E^*$ and hence E is real.

- **Operators** that correspond to **observables** are called **Hermitian operators**.

Hermitian (self-adjoint) operator

- ▶ In QM, **observables** correspond to **operators** that have certain special properties.

A. the expectation value of an **observable** must be a **real number**

$$\langle \hat{O} \rangle = \int \psi^* \hat{O} \psi dx = \langle \psi | \hat{O} \psi \rangle = \langle \hat{O} \rangle^* \quad (1)$$

B. for **operators** that represent **observables**

$$\langle \psi, \hat{O} \psi \rangle = \langle \hat{O} \psi, \psi \rangle \quad \left(\langle g, f \rangle = \langle f, g \rangle^*, \psi(x) \in \mathcal{H} \right) \quad (2)$$

$$\text{Proof :} \quad \hat{O} \psi = E \psi, \quad (3)$$

$$\langle f, \hat{O} f \rangle = \langle f, \hat{O} f \rangle^* = \langle \hat{O} f, f \rangle, \quad (4)$$

$$E \langle f, f \rangle = E^* \langle f, f \rangle \quad (5)$$

where $\langle f, f \rangle$ cannot be zero, so $E = E^*$ and hence E is real.

- ▶ **Operators** that correspond to **observables** are called **Hermitian operators**.
- ▶ self-adjoint :

$$\hat{O} = \hat{O}^\dagger \quad (6)$$

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x)$$

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x) \quad \rightarrow \quad Hu_E(x) = Eu_E(x)$$

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x) \quad \rightarrow \quad Hu_E(x) = Eu_E(x)$$

where E and $u_E(x)$ are eigenvalues and eigenfunctions

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x) \quad \rightarrow \quad Hu_E(x) = Eu_E(x)$$

where E and $u_E(x)$ are eigenvalues and eigenfunctions

- ▶ eigenfunctions for different eigenvalues are orthogonal

$$\int_{-\infty}^{\infty} dx u_{E_1}^*(x) u_{E_2}(x) = 0 \quad \text{where } E_1 \neq E_2 \quad (7)$$

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x) \quad \rightarrow \quad Hu_E(x) = Eu_E(x)$$

where E and $u_E(x)$ are eigenvalues and eigenfunctions

- ▶ eigenfunctions for different eigenvalues are orthogonal

$$\int_{-\infty}^{\infty} dx u_{E_1}^*(x) u_{E_2}(x) = 0 \quad \text{where } E_1 \neq E_2 \quad (7)$$

- ▶ the eigenfunctions become normalized

$$\int_{-\infty}^{\infty} dx u_n^*(x) u_m(x) = \delta_{nm} \quad (8)$$

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x) \quad \rightarrow \quad Hu_E(x) = Eu_E(x)$$

where E and $u_E(x)$ are eigenvalues and eigenfunctions

- ▶ eigenfunctions for different eigenvalues are orthogonal

$$\int_{-\infty}^{\infty} dx u_{E_1}^*(x) u_{E_2}(x) = 0 \quad \text{where } E_1 \neq E_2 \quad (7)$$

- ▶ the eigenfunctions become normalized

$$\int_{-\infty}^{\infty} dx u_n^*(x) u_m(x) = \delta_{nm} \quad (8)$$

- ▶ the eigenfunctions of H form a complete set

$$\psi(x, t) = \sum_n C_n u_n(x) e^{-iE_n t/\hbar} + \int dE C(E) u_E(x) e^{-iEt/\hbar} \quad (9)$$

the C_n and $C(E)$ are arbitrary constants and functions of E .

eigenvalues and eigenfunctions : E and u_E

$$-\frac{\hbar^2}{2m} \frac{d^2 u(x)}{dx^2} + V(x)u(x) = Eu(x) \quad \rightarrow \quad Hu_E(x) = Eu_E(x)$$

where E and $u_E(x)$ are eigenvalues and eigenfunctions

- ▶ eigenfunctions for different eigenvalues are orthogonal

$$\int_{-\infty}^{\infty} dx u_{E_1}^*(x) u_{E_2}(x) = 0 \quad \text{where } E_1 \neq E_2 \quad (7)$$

- ▶ the eigenfunctions become normalized

$$\int_{-\infty}^{\infty} dx u_n^*(x) u_m(x) = \delta_{nm} \quad (8)$$

- ▶ the eigenfunctions of H form a complete set

$$\psi(x, t) = \sum_n C_n u_n(x) e^{-iE_n t/\hbar} + \int dE C(E) u_E(x) e^{-iEt/\hbar} \quad (9)$$

the C_n and $C(E)$ are arbitrary constants and functions of E .

- ▶ the set of all square-integrable functions, on a specific interval,

$$u(x) \text{ such that } \int_a^b |u(x)|^2 dx < \infty \quad (10)$$

$Hu_E(x) = Eu_E(x) : \text{Hilbert space}$

- ▶ the collection of all functions of $u(x)$ constitutes a vector space.

$Hu_E(x) = Eu_E(x) : \text{Hilbert space}$

- ▶ the collection of all functions of $u(x)$ constitutes a vector space.
- ▶ Mathematicians call it $L^2(a, b)$, physicists call it Hilbert space.

$Hu_E(x) = Eu_E(x) : \text{Hilbert space}$

- ▶ the collection of all functions of $u(x)$ constitutes a vector space.
- ▶ Mathematicians call it $L^2(a, b)$, physicists call it Hilbert space.
- ▶ inner product

$$\langle f|g \rangle \equiv \int_a^b f^*(x)g(x)dx = \text{scalar (a real or complex number)}$$

$Hu_E(x) = Eu_E(x)$: Hilbert space

- ▶ the collection of all functions of $u(x)$ constitutes a vector space.
- ▶ Mathematicians call it $L^2(a, b)$, physicists call it Hilbert space.
- ▶ inner product

$$\langle f|g\rangle \equiv \int_a^b f^*(x)g(x)dx = \text{scalar (a real or complex number)}$$

- ▶ square-integrable (norm)

$$\|f\| = \sqrt{\langle f|f\rangle} = \text{positive-definite (non-negative real number)}$$

$Hu_E(x) = Eu_E(x)$: Hilbert space

- ▶ the collection of all functions of $u(x)$ constitutes a vector space.
- ▶ Mathematicians call it $L^2(a, b)$, physicists call it Hilbert space.
- ▶ inner product

$$\langle f|g\rangle \equiv \int_a^b f^*(x)g(x)dx = \text{scalar (a real or complex number)}$$


- ▶ square-integrable (norm)

$$||f|| = \sqrt{\langle f|f\rangle} = \text{positive-definite (non-negative real number)}$$

- ▶ which satisfies completeness

$$f(x) = \sum_{n=1}^{\infty} c_n f_n(x)$$

A particle in a box (an infinite potential well)



$$V(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq L, \\ \infty & \text{otherwise} \end{cases}$$

A particle in a box (an infinite potential well)

$$V(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq L, \\ \infty & \text{otherwise} \end{cases} \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

A particle in a box (an infinite potential well)

$$V(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq L, \\ \infty & \text{otherwise} \end{cases} \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

- ▶ the region outside the box : $\psi(x) = 0$

A particle in a box (an infinite potential well)

$$V(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq L, \\ \infty & \text{otherwise} \end{cases} \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

- ▶ the region outside the box : $\psi(x) = 0$
- ▶ the region inside the box :

$$(\text{B.C.}) \quad \psi(0) = \psi(L) = 0,$$

$$\psi(x) = A \sin(kx) + B \cos(kx) \quad \rightarrow \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)$$

A particle in a box (an infinite potential well)

$$V(x) = \begin{cases} 0 & \text{for } 0 \leq x \leq L, \\ \infty & \text{otherwise} \end{cases} \quad -\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

- ▶ the region outside the box : $\psi(x) = 0$
- ▶ the region inside the box :




$$(\text{B.C.}) \quad \psi(0) = \psi(L) = 0,$$

$$\psi(x) = A \sin(kx) + B \cos(kx) \quad \rightarrow \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)$$

- ▶ the corresponding energy is

$$E_n = \frac{\hbar^2 k^2}{2m} = \frac{n^2 \hbar^2}{8mL^2}$$

A particle in a box : A solution



A particle in a box

► Eigenvalues and Eigenfunctions

$$E_n = \frac{\hbar^2 k^2}{2m} = \frac{n^2 h^2}{8mL^2} = \text{real values}, \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right) \quad (11)$$

A particle in a box

► Eigenvalues and Eigenfunctions

$$E_n = \frac{\hbar^2 k^2}{2m} = \frac{n^2 h^2}{8mL^2} = \text{real values}, \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right) \quad (11)$$

► orthonormal :

$$\int_0^L \psi_n^*(x) \psi_m(x) dx = \frac{2}{L} \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx = \delta_{mn}, \quad (12)$$

A particle in a box

- Eigenvalues and Eigenfunctions

$$E_n = \frac{\hbar^2 k^2}{2m} = \frac{n^2 \hbar^2}{8mL^2} = \text{real values}, \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right) \quad (11)$$

- orthonormal :

$$\int_0^L \psi_n^*(x) \psi_m(x) dx = \frac{2}{L} \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx = \delta_{mn}, \quad (12)$$

- complete set :

$$\psi(x) = \sum_n^{\infty} c_n \psi_n(x) \quad (13)$$

A particle in a box

► Eigenvalues and Eigenfunctions

$$E_n = \frac{\hbar^2 k^2}{2m} = \frac{n^2 \hbar^2}{8mL^2} = \text{real values}, \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right) \quad (11)$$

► orthonormal :

$$\int_0^L \psi_n^*(x) \psi_m(x) dx = \frac{2}{L} \int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx = \delta_{mn}, \quad (12)$$

► complete set :

$$\psi(x) = \sum_n^{\infty} c_n \psi_n(x) \quad (13)$$

Hermitian operator

$$\hat{O} = -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) = \hat{O}^\dagger \quad (14)$$

Summary

In QM,

Summary

In QM,

- ▶ solve Schrödinger equation

Summary

In QM,

- ▶ solve Schrödinger equation
- ▶ Hermitian system

Summary

In QM,

- ▶ solve Schrödinger equation
- ▶ Hermitian system
- ▶ energy conservation

Summary

In QM,

- ▶ solve Schrödinger equation
- ▶ Hermitian system
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)

Summary

In QM,

- ▶ solve Schrödinger equation
- ▶ Hermitian system
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal

Summary

In QM,

- ▶ solve Schrödinger equation
- ▶ Hermitian system
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set

Summary

In QM,

- ▶ solve Schrödinger equation
- ▶ Hermitian system
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

A particle in a black hole

Our background spacetime

Schwarzschild black hole

$$ds^2 = -\left(1 - \frac{2M}{r}\right)dt^2 + \frac{1}{\left(1 - \frac{2M}{r}\right)}dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (15)$$

Our background spacetime

Schwarzschild black hole

$$ds^2 = -\left(1 - \frac{2M}{r}\right)dt^2 + \frac{1}{\left(1 - \frac{2M}{r}\right)}dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (15)$$

- asymptotically flat in 4-dimensional spacetime

Our background spacetime

Schwarzschild black hole

$$ds^2 = -\left(1 - \frac{2M}{r}\right)dt^2 + \frac{1}{\left(1 - \frac{2M}{r}\right)}dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (15)$$

- ▶ asymptotically flat in 4-dimensional spacetime
- ▶ a static solution

Our background spacetime

Schwarzschild black hole

$$ds^2 = -\left(1 - \frac{2M}{r}\right)dt^2 + \frac{1}{\left(1 - \frac{2M}{r}\right)}dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (15)$$

- ▶ asymptotically flat in 4-dimensional spacetime
- ▶ a static solution
- ▶ a vacuum solution $T_{\mu\nu} = 0$

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi T_{\mu\nu} \quad (16)$$

from Schrödinger equation to Klein-Gordon equation

- ▶ Schrödinger eq. : the non-relativistic energy-momentum relation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} \quad \left(\psi = e^{-i(Et - Px)/\hbar} \right) \rightarrow E = \frac{P^2}{2m} \quad (17)$$

- ▶ the relativistic energy-momentum relation :

$$E^2 = P^2 c^2 + m^2 c^4 \quad (18)$$

requires

$$\hat{E} = i\hbar \frac{\partial}{\partial t}, \hat{P} = \frac{\hbar}{i} \frac{\partial}{\partial \vec{x}} \rightarrow -\hbar^2 \frac{\partial^2}{\partial t^2} \psi = -\hbar^2 c^2 \frac{\partial^2}{\partial \vec{x}^2} \psi + m^2 c^4 \psi \quad (19)$$

$$\frac{1}{c^2} \frac{\partial^2}{\partial t^2} \psi - \frac{\partial^2}{\partial \vec{x}^2} \psi + \frac{m^2 c^2}{\hbar^2} \psi = 0 \rightarrow \nabla^2 \psi + m^2 \psi = 0 \quad (20)$$

$$\text{where} \quad \nabla_\mu = \frac{\partial}{\partial x^\mu}, \quad x^\mu = \{ct, x, y, z\}, \quad (21)$$

$$\nabla^2 = \eta^{\mu\nu} \nabla_\mu \nabla_\nu, \quad \eta_{\mu\nu} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \eta^{\mu\nu} \quad (22)$$

Klein-Gordon equation

Quantum mechanics + Relativity $\rightarrow$ Quantum field theory

Klein-Gordon equation: $\nabla^2 \psi + m^2 \psi = 0$ (23)

- ▶ ψ : fields (not a wave function)
- ▶ $\langle \psi | \psi \rangle = \int d^3x |\psi(x)|^2$: not Lorentz invariant
- ▶ Klein-Gordon inner product :

$$\langle \psi | \psi \rangle_{\text{KG}} = \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \quad (24)$$

which is invariant under Lorentz transformation and ensures conservation under time evolution

Klein-Gordon equation on curved spacetime (I)

- Klein-Gordon equation on curved spacetime ($m^2 = 0$)

$$\nabla^2 \psi = \nabla^\mu \nabla_\mu \psi = \nabla_\mu g^{\mu\nu} \nabla_\nu \psi = 0, \quad (\nabla_\mu g_{\alpha\beta} = 0) \quad (25)$$

$$g^{\mu\nu} \nabla_\mu (\partial_\nu \psi) = g^{\mu\nu} \left(\partial_\mu (\partial_\nu \psi) - \Gamma_{\mu\nu}^\gamma (\partial_\gamma \psi) \right) = 0 \quad (26)$$

where $\nabla_\mu \psi = \partial_\mu \psi$ and $\nabla_\mu A_\nu = \partial_\mu A_\nu - \Gamma_{\mu\nu}^\gamma A_\gamma$

and

$$ds^2 = -A(r)dt^2 + \frac{1}{A(r)}dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (27)$$

Klein-Gordon equation on curved spacetime (I)

Klein-Gordon equation on curved spacetime ($m^2 = 0$)

$$\nabla^2 \psi = g^{\mu\nu} \left(\partial_\mu (\partial_\nu \psi) - \Gamma_{\mu\nu}^\gamma (\partial_\gamma \psi) \right) = 0 \quad (28)$$

$$ds^2 = -A(r)dt^2 + \frac{1}{A(r)}dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \quad (29)$$

$$g_{\mu\nu} = \begin{pmatrix} -A(r) & 0 & 0 & 0 \\ 0 & \frac{1}{A(r)} & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2 \theta \end{pmatrix}, \quad g_{\mu\nu} g^{\nu\gamma} = \delta_\mu^\gamma \quad (30)$$

$$\Gamma_{\mu\nu}^\gamma = \frac{1}{2} g^{\gamma\delta} \left[\frac{\partial g_{\mu\delta}}{\partial x^\nu} + \frac{\partial g_{\nu\delta}}{\partial x^\mu} - \frac{\partial g_{\mu\nu}}{\partial x^\delta} \right], \quad (31)$$

$$\Gamma_{tt}^t = \frac{1}{2} g^{t\delta} \left[\frac{\partial g_{t\delta}}{\partial x^t} + \frac{\partial g_{t\delta}}{\partial x^t} - \frac{\partial g_{tt}}{\partial x^\delta} \right] = \frac{1}{2} g^{tt} \left[\frac{\partial g_{tt}}{\partial x^t} + \frac{\partial g_{tt}}{\partial x^t} - \frac{\partial g_{tt}}{\partial x^t} \right] = 0$$

$$\begin{aligned} \Gamma_{tr}^t &= \frac{1}{2} g^{t\delta} \left[\frac{\partial g_{t\delta}}{\partial x^r} + \frac{\partial g_{r\delta}}{\partial x^t} - \frac{\partial g_{tr}}{\partial x^\delta} \right] = \frac{1}{2} g^{tt} \left[\frac{\partial g_{tt}}{\partial x^r} + \frac{\partial g_{tt}}{\partial x^t} - \frac{\partial g_{tr}}{\partial x^t} \right] \\ &= \frac{1}{2} \left(-\frac{1}{A(r)} \right) \left[\frac{\partial(-A(r))}{\partial x^r} + \frac{\partial(-A(r))}{\partial x^t} - 0 \right] = \frac{A'(r)}{2A(r)} \end{aligned} \quad (32)$$

Klein-Gordon equation on curved spacetime (I)

$$\Gamma_{rt}^t = \frac{A'(r)}{2A(r)}, \quad \Gamma_{tt}^r = \frac{1}{2}A(r)A'(r), \quad \Gamma_{rr}^r = -\frac{A'(r)}{2A(r)}, \quad (28)$$

$$\Gamma_{\theta\theta}^r = -rA(r), \quad \Gamma_{\phi\phi}^r = -rA(r)\sin^2(\theta), \quad \Gamma_{\theta r}^\theta = \frac{1}{r}, \quad (29)$$

$$\Gamma_{\phi\phi}^\theta = -\cos(\theta)\sin(\theta), \quad \Gamma_{\phi r}^\phi = \frac{1}{r}, \quad \Gamma_{\phi\theta}^\phi = \cot(\theta) \quad (30)$$

$$\begin{aligned} \nabla^2\psi &= g^{\mu\nu} \left(\partial_\mu(\partial_\nu\psi) - \Gamma_{\mu\nu}^\gamma(\partial_\gamma\psi) \right) \\ &= g^{tt} \left(\partial_t(\partial_t\psi) - \Gamma_{tt}^\gamma(\partial_\gamma\psi) \right) + g^{rr} \left(\partial_r(\partial_r\psi) - \Gamma_{rr}^\gamma(\partial_\gamma\psi) \right) \\ &\quad + g^{\theta\theta} \left(\partial_\theta(\partial_\theta\psi) - \Gamma_{\theta\theta}^\gamma(\partial_\gamma\psi) \right) + g^{\phi\phi} \left(\partial_\phi(\partial_\phi\psi) - \Gamma_{\phi\phi}^\gamma(\partial_\gamma\psi) \right) \\ &= g^{tt} \left(\partial_t(\partial_t\psi) - \Gamma_{tt}^r(\partial_r\psi) \right) + g^{rr} \left(\partial_r(\partial_r\psi) - \Gamma_{rr}^r(\partial_r\psi) \right) \\ &\quad + g^{\theta\theta} \left(\partial_\theta(\partial_\theta\psi) - \Gamma_{\theta\theta}^r(\partial_r\psi) \right) + g^{\phi\phi} \left(\partial_\phi(\partial_\phi\psi) - \Gamma_{\phi\phi}^r(\partial_r\psi) - \Gamma_{\phi\phi}^\theta(\partial_\theta\psi) \right) \end{aligned}$$

Klein-Gordon equation on curved spacetime (I)

$$\psi(t, r, \theta, \phi) = e^{-i\omega t} \frac{\Phi(r)}{r} Y_m^l(\theta, \phi), \quad (31)$$

$$\begin{aligned} \nabla^2 \psi = & g^{tt} \left[\partial_t (\partial_t \psi) - \frac{A(r)A'(r)}{2} (\partial_r \psi) \right] \\ & + g^{rr} \left[\partial_r (\partial_r \psi) - \left(-\frac{A'(r)}{2A(r)} \right) (\partial_r \psi) \right] + g^{\theta\theta} \left[\partial_\theta (\partial_\theta \psi) - (-rA(r)) (\partial_r \psi) \right] \\ & + g^{\phi\phi} \left[\partial_\phi (\partial_\phi \psi) - (-rA(r) \sin^2 \theta) (\partial_r \psi) - (-\cos \theta \sin \theta) (\partial_\theta \psi) \right] \end{aligned} \quad (32)$$

using

$$\frac{1}{Y_m^l} \frac{\partial^2 Y_m^l}{\partial \phi^2} = -m^2, \quad l(l+1) \sin^2 \theta + \frac{\sin \theta}{Y_m^l} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial Y_m^l}{\partial \theta} \right) = m^2 \quad (33)$$

then

$$\Phi''(r) + \frac{A'(r)}{A(r)} \Phi'(r) - \frac{1}{A(r)} \left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} - \frac{\omega^2}{A(r)} \right) \Phi(r) = 0 \quad (34)$$

Klein-Gordon equation on curved spacetime (II)

$$\nabla^2 \psi(t, r, \theta, \phi) = \frac{1}{\sqrt{-g}} \partial_\mu (g^{\mu\nu} \sqrt{-g} \partial_\nu \psi) = 0, \quad (35)$$

$$g_{\mu\nu} = \begin{pmatrix} -A(r) & 0 & 0 & 0 \\ 0 & \frac{1}{A(r)} & 0 & 0 \\ 0 & 0 & r^2 & 0 \\ 0 & 0 & 0 & r^2 \sin^2(\theta) \end{pmatrix}, \quad \sqrt{-g} = r^2 \sin(\theta) \quad (36)$$

$$g^{\mu\nu} = \begin{pmatrix} -\frac{1}{A(r)} & 0 & 0 & 0 \\ 0 & A(r) & 0 & 0 \\ 0 & 0 & \frac{1}{r^2} & 0 \\ 0 & 0 & 0 & \frac{1}{r^2 \sin^2(\theta)} \end{pmatrix} \quad (37)$$

$$\psi(t, r, \theta, \phi) = e^{-i\omega t} \frac{\Phi(r)}{r} Y_m^l(\theta, \phi) \quad (38)$$

$$\begin{aligned} \nabla^2 \psi(t, r, \theta, \phi) = & \frac{1}{r^2 \sin(\theta)} \left[\partial_t \left(g^{tt} r^2 \sin(\theta) \partial_t \psi \right) + \partial_r \left(g^{rr} r^2 \sin(\theta) \partial_r \psi \right) \right. \\ & \left. + \partial_\theta \left(g^{\theta\theta} r^2 \sin(\theta) \partial_\theta \psi \right) + \partial_\phi \left(g^{\phi\phi} r^2 \sin(\theta) \partial_\phi \psi \right) \right] = 0 \end{aligned} \quad (39)$$

Klein-Gordon equation on curved spacetime (II)

$$\begin{aligned} \nabla^2 \psi(t, r, \theta, \phi) = & \partial_t \left(-\frac{1}{A(r)} \partial_t \psi \right) + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \psi \right) \\ & + \frac{1}{r^2 \sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta \psi \right) + \frac{1}{r^2 \sin^2(\theta)} \partial_\phi \left(\partial_\phi \psi \right) \end{aligned} \quad (40)$$

Let us insert : $\psi(t, r, \theta, \phi) = e^{-i\omega t} \frac{\Phi(r)}{r} Y_m^l(\theta, \phi)$

$$= \partial_t \left(-\frac{1}{A(r)} \partial_t e^{-i\omega t} \right) \frac{\Phi(r)}{r} Y_m^l + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \frac{\Phi(r)}{r} \right) e^{-i\omega t} Y_m^l$$

Klein-Gordon equation on curved spacetime (II)

$$\begin{aligned} \nabla^2 \psi(t, r, \theta, \phi) = & \partial_t \left(-\frac{1}{A(r)} \partial_t \psi \right) + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \psi \right) \\ & + \frac{1}{r^2 \sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta \psi \right) + \frac{1}{r^2 \sin^2(\theta)} \partial_\phi \left(\partial_\phi \psi \right) \end{aligned} \quad (40)$$

Let us insert : $\psi(t, r, \theta, \phi) = e^{-i\omega t} \frac{\Phi(r)}{r} Y_m^l(\theta, \phi)$

$$\begin{aligned} = & \partial_t \left(-\frac{1}{A(r)} \partial_t e^{-i\omega t} \right) \frac{\Phi(r)}{r} Y_m^l + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \frac{\Phi(r)}{r} \right) e^{-i\omega t} Y_m^l \\ & + \frac{1}{r^2 \sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta Y_m^l \right) e^{-i\omega t} \frac{\Phi(r)}{r} + \frac{1}{r^2 \sin^2(\theta)} \partial_\phi \left(\partial_\phi Y_m^l \right) e^{-i\omega t} \frac{\Phi(r)}{r} \end{aligned}$$

Klein-Gordon equation on curved spacetime (II)

$$\begin{aligned}\nabla^2\psi(t, r, \theta, \phi) = & \partial_t \left(-\frac{1}{A(r)} \partial_t \psi \right) + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \psi \right) \\ & + \frac{1}{r^2 \sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta \psi \right) + \frac{1}{r^2 \sin^2(\theta)} \partial_\phi \left(\partial_\phi \psi \right)\end{aligned}\quad (40)$$

Let us insert : $\psi(t, r, \theta, \phi) = e^{-i\omega t} \frac{\Phi(r)}{r} Y_m^l(\theta, \phi)$

$$\begin{aligned}= & \partial_t \left(-\frac{1}{A(r)} \partial_t e^{-i\omega t} \right) \frac{\Phi(r)}{r} Y_m^l + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \frac{\Phi(r)}{r} \right) e^{-i\omega t} Y_m^l \\ & + \frac{1}{r^2} \left[\frac{1}{\sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta Y_m^l \right) + \frac{1}{\sin^2(\theta)} \partial_\phi \left(\partial_\phi Y_m^l \right) \right] e^{-i\omega t} \frac{\Phi(r)}{r}\end{aligned}\quad (41)$$

$$\left(\frac{1}{\sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta Y_m^l \right) + \frac{1}{\sin^2(\theta)} \partial_\phi \left(\partial_\phi Y_m^l \right) = -l(l+1) Y_m^l \right) \quad (42)$$

Klein-Gordon equation on curved spacetime (II)

$$\begin{aligned}\nabla^2\psi(t, r, \theta, \phi) &= \partial_t \left(-\frac{1}{A(r)} \partial_t \psi \right) + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \psi \right) \\ &+ \frac{1}{r^2 \sin(\theta)} \partial_\theta \left(\sin(\theta) \partial_\theta \psi \right) + \frac{1}{r^2 \sin^2(\theta)} \partial_\phi \left(\partial_\phi \psi \right)\end{aligned}\quad (40)$$

Let us insert : $\psi(t, r, \theta, \phi) = e^{-i\omega t} \frac{\Phi(r)}{r} Y_m^l(\theta, \phi)$

$$\begin{aligned}&= \partial_t \left(-\frac{1}{A(r)} \partial_t e^{-i\omega t} \right) \frac{\Phi(r)}{r} Y_m^l + \frac{1}{r^2} \partial_r \left(A(r) r^2 \partial_r \frac{\Phi(r)}{r} \right) e^{-i\omega t} Y_m^l \\ &+ \frac{1}{r^2} \left[-l(l+1) Y_m^l \right] e^{-i\omega t} \frac{\Phi(r)}{r}\end{aligned}\quad (41)$$

Klein-Gordon equation on curved spacetime (II)

$$\begin{aligned}\nabla^2\psi(t,r,\theta,\phi) &= \partial_t\left(-\frac{1}{A(r)}\partial_t\psi\right) + \frac{1}{r^2}\partial_r\left(A(r)r^2\partial_r\psi\right) \\ &+ \frac{1}{r^2\sin(\theta)}\partial_\theta\left(\sin(\theta)\partial_\theta\psi\right) + \frac{1}{r^2\sin^2(\theta)}\partial_\phi\left(\partial_\phi\psi\right)\end{aligned}\quad (40)$$

Let us insert : $\psi(t,r,\theta,\phi) = e^{-i\omega t}\frac{\Phi(r)}{r}Y_m^l(\theta,\phi)$

$$\begin{aligned}&= \partial_t\left(-\frac{1}{A(r)}\partial_t e^{-i\omega t}\right)\frac{\Phi(r)}{r}Y_m^l + \frac{1}{r^2}\partial_r\left(A(r)r^2\partial_r\frac{\Phi(r)}{r}\right)e^{-i\omega t}Y_m^l \\ &+ \frac{1}{r^2}\left[-l(l+1)Y_m^l\right]e^{-i\omega t}\frac{\Phi(r)}{r}\end{aligned}\quad (41)$$

$$\rightarrow \Phi''(r) + \frac{A'(r)}{A(r)}\Phi'(r) - \frac{1}{A(r)}\left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} - \frac{\omega^2}{A(r)}\right)\Phi(r) = 0$$

Klein-Gordon equation on curved spacetime

A scalar field equation on curved spacetime

$$\Phi''(r) + \frac{A'(r)}{A(r)}\Phi'(r) - \frac{1}{A(r)}\left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} - \frac{\omega^2}{A(r)}\right)\Phi(r) = 0 \quad (42)$$

Let us employ a tortoise coordinate

$$dr_* = \frac{1}{A(r)}dr \quad \left(ds^2 = -A(r)(dt^2 + dr_*^2) + r^2 d\Omega_2\right) \quad (43)$$

it becomes Schrodinger equation like

$$\frac{d^2}{dr_*^2}\Phi''(r) - V_{\text{eff}}\Phi(r) = -\omega^2\Phi(r) \quad (44)$$

where

$$V_{\text{eff}} = A(r)\left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r}\right) \quad (45)$$

We will solve this equation classically (not quantize).

Quasi-Normal Modes (QNM)

$$\frac{d^2}{dr_*^2} \Phi''(r) + V_{\text{eff}} \Phi(r) = -\omega^2 \Phi(r) \quad (46)$$

Boundary conditions

- ▶ near the horizon : incoming, $e^{-i(t+r)\omega}$
- ▶ at infinity : outgoing, $e^{-i(t-r)\omega}$

which indicates "*open system*"

- ▶ non-Hermitian : $\mathcal{L} \neq \mathcal{L}^\dagger$
- ▶ no conservation in time evolution
- ▶ the Klein-Gordon inner product at infinity diverge
- ▶ ω is not real but complex (discrete and infinity number of set)
→ Quasi-Normal Modes (QNM)

How to impose the Boundary Conditions?

$$\frac{d^2}{dr_*^2} \Phi''(r) + V_{\text{eff}} \Phi(r) = -\omega^2 \Phi(r) \quad (47)$$

- ▶ (B.C.) near the horizon : incoming, $e^{-i(t+r_*)\omega}$
- ▶ (B.C.) at infinity : outgoing, $e^{-i(t-r_*)\omega}$

$$\begin{aligned} V_{\text{eff}} = 0 : \quad \Phi(r) &= c_1 \cos(r_* \omega) + c_2 \sin(r_* \omega) \\ &= \frac{1}{2} c_1 e^{-ir_* \omega} + \frac{1}{2} c_1 e^{ir_* \omega} - \frac{1}{2} i c_2 e^{ir_* \omega} + \frac{1}{2} i c_2 e^{-ir_* \omega} \end{aligned} \quad (48)$$

- incoming boundary condition : $c_1 = i c_2$

$$\Phi(r) = i c_2 e^{-ir_* \omega}, \quad e^{-i\omega t} \Phi(r) = -i c_2 e^{-i(t+r_*)\omega} \quad (49)$$

- outgoing boundary condition : $c_1 = -i c_2$


$$\Phi(r) = -i c_2 e^{ir_* \omega}, \quad e^{-i\omega t} \Phi(r) = -i c_2 e^{-i(t-r_*)\omega} \quad (50)$$

- ▶ previously (a particle in a box)


$$(B.C.) \quad V_{\text{eff}} = 0 \quad (0 \leq x \leq L), \quad \psi(0) = \psi(L) = 0,$$

$$\psi(x) = A \sin(kx) + B \cos(kx) \quad \rightarrow \quad \psi_n(x) = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi x}{L}\right)$$


A particle in a box



A particle in a black hole



V_{eff} : effective potential for KG field



How to obtain QNMs (ω) : 1. Eigenvalues Problem

$$\frac{d^2}{dr_*^2} \Phi''(r) + V_{\text{eff}} \Phi(r) = -\omega^2 \Phi(r) \quad (51)$$

which takes a form of

$$\mathcal{L} \Phi(r) = \lambda \Phi(r) \quad (52)$$


this is nothing but the eigenvalue problem.

$$\mathcal{L} u_n = \lambda_n u_n \quad (53)$$

- By using numerical methods (e.g. pseudo spectral methods or shooting method), we impose the boundary condition for QNM and can calculate λ_n and u_n .

How to obtain QNMs (ω) : 1. Eigenvalues Problem

QNM with $l = 0$ for scalar field in Schwarzschild black hole by using pseudo spectral method





	$\text{Re}[\omega_n]$	$\text{Im}[\omega_n]$
1	± 0.13024640	-2.7140490
2	± 0.13444262	-2.2112523
3	± 0.14078647	-1.7073040
4	± 0.15148323	-1.2021605
5	± 0.17223383	-0.69610489
6	± 0.22090988	-0.20979143

$$e^{-i\omega t} = e^{-i(\omega_R + i\omega_I)t} = e^{-i\omega_R t} e^{\omega_I t}$$

How to obtain QNMs (ω) : 1. Eigenvalues Problem


QNM with $l = 0$ for scalar field in Schwarzschild black hole by using pseudo spectral method



$$e^{-i\omega t} = e^{-i(\omega_R + i\omega_I)t} = e^{-i\omega_R t} e^{\omega_I t}$$


How to obtain QNMs (ω) : 1. Eigenvalues Problem

QNM with $l = 0, 1, 2, 3$ for scalar field in Schwarzschild black hole by using pseudo spectral method



How to obtain QNMs (ω) : 1. Eigenvalues Problem

QNM with $l = 0$ for scalar field in Schwarzschild black hole by using pseudo spectral method in time domain



How to obtain QNMs (ω) : 2. WKB method ¹

$$\frac{d^2}{dr_*^2} \Phi''(r) + V_{\text{eff}} \Phi(r) = -\omega^2 \Phi(r)$$




FIG. 1. The function $V(r_*) - \omega^2$.

¹Sai Iyer, "Black-hole normal modes: A WKB approach. II. Schwarzschild black holes", Phys. Rev. D 35, 3632

How to obtain QNMs (ω) : 2. WKB method ¹

$$\frac{d^2}{dr_*^2} \Phi''(r) + V_{\text{eff}} \Phi(r) = -\omega^2 \Phi(r)$$




FIG. 1. The function $V(r_*) - \omega^2$.

mula for the normal-mode frequencies, given by

$$\omega^2 = [V_0 + (-2V_0'')^{1/2} \tilde{\Lambda}]$$

$$-i(n + \frac{1}{2})(-2V_0'')^{1/2}(1 + \tilde{\Omega}),$$

(1.2) where

$$n = \begin{cases} 0, 1, 2, \dots, & \text{Re}(\omega) > 0, \\ -1, -2, -3, \dots, & \text{Re}(\omega) < 0, \end{cases}$$

$$\tilde{\Lambda}(n) = \frac{1}{(-2V_0'')^{1/2}} \left[\frac{1}{8} \left(\frac{V_0^{(4)}}{V_0''} \right) \left[\frac{1}{4} + \alpha^2 \right] - \frac{1}{288} \left(\frac{V_0'''}{V_0''} \right)^2 (7 + 60\alpha^2) \right], \quad (1.3a)$$

$$\tilde{\Omega}(n) = \frac{1}{(-2V_0'')^{1/2}} \left[\frac{5}{6912} \left(\frac{V_0'''}{V_0''} \right)^4 (77 + 188\alpha^2) - \frac{1}{384} \left(\frac{V_0'''' V_0^{(4)}}{V_0''^3} \right) (51 + 100\alpha^2) + \frac{1}{2304} \left(\frac{V_0^{(4)}}{V_0''} \right)^2 (67 + 68\alpha^2) \right. \\ \left. + \frac{1}{288} \left(\frac{V_0'''' V_0^{(5)}}{V_0''^2} \right) (19 + 28\alpha^2) - \frac{1}{288} \left(\frac{V_0^{(6)}}{V_0''} \right) (5 + 4\alpha^2) \right]. \quad (1.3b)$$

¹Sai Iyer, "Black-hole normal modes: A WKB approach. II. Schwarzschild black holes", Phys. Rev. D 35, 3632

How to obtain QNMs (ω) : 2. WKB method ¹

$$\frac{d^2}{dr_*^2} \Phi''(r) + V_{\text{eff}} \Phi(r) = -\omega^2 \Phi(r)$$




FIG. 1. The function $V(r_*) - \omega^2$.

mula for the normal-mode frequencies, given by

$$\omega^2 = [V_0 + (-2V_0'')^{1/2} \tilde{\Lambda}] - i(n + \frac{1}{2})(-2V_0'')^{1/2}(1 + \tilde{\Omega}), \quad (1.2)$$

where

$$\tilde{\Lambda}(n) = \frac{1}{(-2V_0'')^{1/2}} \left[\frac{1}{8} \left[\frac{V_0^{(4)}}{V_0''} \right] \left[\frac{1}{4} + \alpha^2 \right] - \frac{1}{288} \left[\frac{V_0'''}{V_0''} \right]^2 (7 + 60\alpha^2) \right], \quad (1.3a)$$

$$\tilde{\Omega}(n) = \frac{1}{(-2V_0'')^{1/2}} \left[\frac{5}{6912} \left[\frac{V_0'''}{V_0''} \right]^4 (77 + 188\alpha^2) - \frac{1}{384} \left[\frac{V_0'''' V_0^{(4)}}{V_0''^3} \right] (51 + 100\alpha^2) + \frac{1}{2304} \left[\frac{V_0^{(4)}}{V_0''} \right]^2 (67 + 68\alpha^2) + \frac{1}{288} \left[\frac{V_0'''' V_0^{(5)}}{V_0''^2} \right] (19 + 28\alpha^2) - \frac{1}{288} \left[\frac{V_0^{(6)}}{V_0''} \right] (5 + 4\alpha^2) \right]. \quad (1.3b)$$




FIG. 2. Gravitational normal modes. Here $\sigma = M\omega$, where M is the mass of the black hole.

¹Sai Iyer, "Black-hole normal modes: A WKB approach. II. Schwarzschild black holes", Phys. Rev. D 35, 3632

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial t^2} - V(x) \right] \psi(t, x) = 0, \quad x = \frac{r_*}{c}, \quad (54)$$

$$\text{Laplace transformation : } \hat{\psi}(s, x) = \int_0^\infty dt e^{-st} \psi(t, x), \quad (55)$$

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}(s, x) = \mathcal{J}(s, x), \quad (s = -i\omega) \quad (56)$$

$$\text{where } \mathcal{J}(s, x) = -\partial_t \psi(t, x) - s\psi(t, x) \Big|_{t=0} \quad (57)$$

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{G}(s, x, x') = \delta(x - x') \quad (58)$$

Then the corresponding solution is given by

$$\hat{\psi}(s, x) = \int_{-\infty}^{\infty} dx' \hat{G}(s, x, x') \mathcal{J}(s, x') \quad (59)$$

If the initial data $\mathcal{J}(s, x')$ is given the solution is uniquely determined.

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}_{\pm}(s, x) = 0,$$

where $\hat{\psi}_{\pm}$ are any two linearly independent sol's of homogenous eq.

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}_{\pm}(s, x) = 0,$$

where $\hat{\psi}_{\pm}$ are any two linearly independent sol's of homogenous eq.

$$\hat{G} = \frac{1}{\mathcal{W}(s)} \left[\Theta(x - x') \hat{\psi}_{-}(s, x') \hat{\psi}_{+}(s, x) + \Theta(x' - x) \hat{\psi}_{-}(s, x) \hat{\psi}_{+}(s, x') \right],$$

where $\mathcal{W}(s) = \hat{\psi}_{-}(s, x) \partial_x \hat{\psi}_{+}(s, x) - \hat{\psi}_{+}(s, x) \partial_x \hat{\psi}_{-}(s, x),$

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}_{\pm}(s, x) = 0,$$

where $\hat{\psi}_{\pm}$ are any two linearly independent sol's of homogenous eq.

$$\hat{G} = \frac{1}{\mathcal{W}(s)} \left[\underbrace{\Theta(x - x') \hat{\psi}_{-}(s, x') \hat{\psi}_{+}(s, x)}_{\text{a field propagates from } x' \text{ to } x} + \underbrace{\Theta(x' - x) \hat{\psi}_{-}(s, x) \hat{\psi}_{+}(s, x')}_{\text{a field propagates from } x \text{ to } x'} \right],$$

where $\mathcal{W}(s) = \hat{\psi}_{-}(s, x) \partial_x \hat{\psi}_{+}(s, x) - \hat{\psi}_{+}(s, x) \partial_x \hat{\psi}_{-}(s, x),$

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}_{\pm}(s, x) = 0,$$

where $\hat{\psi}_{\pm}$ are any two linearly independent sol's of homogenous eq.

$$\hat{G} = \frac{1}{\mathcal{W}(s)} \left[\underbrace{\Theta(x - x') \hat{\psi}_{-}(s, x') \hat{\psi}_{+}(s, x)}_{\text{a field propagates from } x' \text{ to } x} + \underbrace{\Theta(x' - x) \hat{\psi}_{-}(s, x) \hat{\psi}_{+}(s, x')}_{\text{a field propagates from } x \text{ to } x'} \right],$$

where $\mathcal{W}(s) = \hat{\psi}_{-}(s, x) \partial_x \hat{\psi}_{+}(s, x) - \hat{\psi}_{+}(s, x) \partial_x \hat{\psi}_{-}(s, x)$,

$$\begin{aligned} \hat{\psi}(s, x) &= \int_{-\infty}^{\infty} dx' \hat{G}(s, x, x') \mathcal{J}(s, x') \\ &= \frac{\hat{\psi}_{+}(s, x)}{\mathcal{W}(s)} \int_{-\infty}^x dx' \hat{\psi}_{-}(s, x') \mathcal{J}(s, x') + \frac{\hat{\psi}_{-}(s, x)}{\mathcal{W}(s)} \int_x^{\infty} dx' \hat{\psi}_{+}(s, x') \mathcal{J}(s, x') \end{aligned}$$

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}_{\pm}(s, x) = 0,$$

where $\hat{\psi}_{\pm}$ are any two linearly independent sol's of homogenous eq.

$$\hat{G} = \frac{1}{\mathcal{W}(s)} \left[\underbrace{\Theta(x - x') \hat{\psi}_{-}(s, x') \hat{\psi}_{+}(s, x)}_{\text{a field propagates from } x' \text{ to } x} + \underbrace{\Theta(x' - x) \hat{\psi}_{-}(s, x) \hat{\psi}_{+}(s, x')}_{\text{a field propagates from } x \text{ to } x'} \right],$$

where $\mathcal{W}(s) = \hat{\psi}_{-}(s, x) \partial_x \hat{\psi}_{+}(s, x) - \hat{\psi}_{+}(s, x) \partial_x \hat{\psi}_{-}(s, x)$,

$$\begin{aligned} \hat{\psi}(s, x) &= \int_{-\infty}^{\infty} dx' \hat{G}(s, x, x') \mathcal{J}(s, x') \\ &= \frac{\hat{\psi}_{+}(s, x)}{\mathcal{W}(s)} \int_{-\infty}^x dx' \hat{\psi}_{-}(s, x') \mathcal{J}(s, x') + \frac{\hat{\psi}_{-}(s, x)}{\mathcal{W}(s)} \int_x^{\infty} dx' \hat{\psi}_{+}(s, x') \mathcal{J}(s, x') \end{aligned}$$

$$\text{If } x > x_R : \hat{\psi}(s, x) = \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x), \quad c_{-}(s) = \int_{x_L}^{x_R} dx' \hat{\psi}_{-}(s, x') \mathcal{J}(s, x')$$

$$\text{If } x < x_L : \hat{\psi}(s, x) = \frac{c_{+}(s)}{\mathcal{W}(s)} \hat{\psi}_{-}(s, x), \quad c_{+}(s) = \int_{x_L}^{x_R} dx' \hat{\psi}_{+}(s, x') \mathcal{J}(s, x')$$

Solution for KG equation: Green's function method

$$\left[\frac{\partial^2}{\partial x^2} - s^2 - V(x) \right] \hat{\psi}_{\pm}(s, x) = 0,$$

where $\hat{\psi}_{\pm}$ are any two linearly independent sol's of homogenous eq.

$$\hat{G} = \frac{1}{\mathcal{W}(s)} \left[\underbrace{\Theta(x - x') \hat{\psi}_{-}(s, x') \hat{\psi}_{+}(s, x)}_{\text{a field propagates from } x' \text{ to } x} + \underbrace{\Theta(x' - x) \hat{\psi}_{-}(s, x) \hat{\psi}_{+}(s, x')}_{\text{a field propagates from } x \text{ to } x'} \right],$$

where $\mathcal{W}(s) = \hat{\psi}_{-}(s, x) \partial_x \hat{\psi}_{+}(s, x) - \hat{\psi}_{+}(s, x) \partial_x \hat{\psi}_{-}(s, x)$,

$$\begin{aligned} \hat{\psi}(s, x) &= \int_{-\infty}^{\infty} dx' \hat{G}(s, x, x') \mathcal{J}(s, x') \\ &= \frac{\hat{\psi}_{+}(s, x)}{\mathcal{W}(s)} \int_{-\infty}^x dx' \hat{\psi}_{-}(s, x') \mathcal{J}(s, x') + \frac{\hat{\psi}_{-}(s, x)}{\mathcal{W}(s)} \int_x^{\infty} dx' \hat{\psi}_{+}(s, x') \mathcal{J}(s, x') \end{aligned}$$

$$\text{If } x > x_R : \hat{\psi}(s, x) = \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x), \quad c_{-}(s) = \int_{x_L}^{x_R} dx' \hat{\psi}_{-}(s, x') \mathcal{J}(s, x')$$

$$\text{If } x < x_L : \hat{\psi}(s, x) = \frac{c_{+}(s)}{\mathcal{W}(s)} \hat{\psi}_{-}(s, x), \quad c_{+}(s) = \int_{x_L}^{x_R} dx' \hat{\psi}_{+}(s, x') \mathcal{J}(s, x')$$

Solution for KG equation: Green's function method

$$\hat{\psi}(s, x) = \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x),$$

Solution for KG equation: Green's function method

$$\hat{\psi}(s, x) = \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x),$$

$$\psi(t, x) = \frac{1}{2\pi i} \int_{\epsilon-i\infty}^{\epsilon+i\infty} ds e^{st} \hat{\psi}(s, x) = \frac{1}{2\pi i} \int_{\epsilon-i\infty}^{\epsilon+i\infty} ds e^{st} \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x)$$


which is known as *Bromwich integral* or *Mellin's inverse formula*.

Solution for KG equation: Green's function method

$$\hat{\psi}(s, x) = \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x),$$

$$\psi(t, x) = \frac{1}{2\pi i} \int_{\epsilon-i\infty}^{\epsilon+i\infty} ds e^{st} \hat{\psi}(s, x) = \frac{1}{2\pi i} \int_{\epsilon-i\infty}^{\epsilon+i\infty} ds e^{st} \frac{c_{-}(s)}{\mathcal{W}(s)} \hat{\psi}_{+}(s, x)$$

which is known as *Bromwich integral* or *Mellin's inverse formula*.




Solution for KG equation: Green's function method

$$\hat{\psi}(s, x) = \frac{c_-(s)}{\mathcal{W}(s)} \hat{\psi}_+(s, x),$$

$$\psi(t, x) = \frac{1}{2\pi i} \int_{\epsilon - i\infty}^{\epsilon + i\infty} ds e^{st} \hat{\psi}(s, x) = \frac{1}{2\pi i} \int_{\epsilon - i\infty}^{\epsilon + i\infty} ds e^{st} \frac{c_-(s)}{\mathcal{W}(s)} \hat{\psi}_+(s, x)$$

which is known as *Bromwich integral* or *Mellin's inverse formula*.



If $\hat{\psi}_+(s, x)$ are analytic in s (there are no essential singularities or branch cuts inside of the contour), the only contribution to the integral comes from the poles of the integrand due to zeros of $\mathcal{W}(s)$.

$$\mathcal{W}(s) = \mathcal{W}'(s_n)(s - s_n) + \mathcal{O}(s - s_n)^2,$$


$$\text{where } \mathcal{W}'(s_n) \equiv \left. \frac{d\mathcal{W}}{ds} \right|_{s=s_n}$$

Solution for KG equation: Green's function method

$$\hat{\psi}(s, x) = \frac{c_-(s)}{\mathcal{W}(s)} \hat{\psi}_+(s, x),$$

$$\psi(t, x) = \frac{1}{2\pi i} \int_{\epsilon - i\infty}^{\epsilon + i\infty} ds e^{st} \hat{\psi}(s, x) = \frac{1}{2\pi i} \int_{\epsilon - i\infty}^{\epsilon + i\infty} ds e^{st} \frac{c_-(s)}{\mathcal{W}(s)} \hat{\psi}_+(s, x)$$

which is known as *Bromwich integral* or *Mellin's inverse formula*.



If $\hat{\psi}_+(s, x)$ are analytic in s (there are no essential singularities or branch cuts inside of the contour), the only contribution to the integral comes from the poles of integrand due to zeros of $\mathcal{W}(s)$.

$$\mathcal{W}(s) = \mathcal{W}'(s_n)(s - s_n) + \mathcal{O}(s - s_n)^2,$$

$$\text{where } \mathcal{W}'(s_n) \equiv \left. \frac{d\mathcal{W}}{ds} \right|_{s=s_n}$$

Solution for KG equation: Green's function method

$$\mathcal{W}(s_n) = \hat{\psi}_-(s_n, x) \partial_x \hat{\psi}_+(s_n, x) - \hat{\psi}_+(s_n, x) \partial_x \hat{\psi}_-(s_n, x) = 0 \quad (60)$$

implies

$$\frac{\hat{\psi}'_+(s_n, x)}{\hat{\psi}_+(s_n, x)} = \frac{\hat{\psi}'_-(s_n, x)}{\hat{\psi}_-(s_n, x)} \quad (61)$$

which integrates to

$$\hat{\psi}_+(s_n, x) = c(s_n) \hat{\psi}_-(s_n, x) \quad (62)$$

"This means that, for $s = s_n$ the two solutions of the homogenous equation, $\hat{\phi}_+(s, x)$ and $\hat{\phi}_-(s, x)$, are no longer independent, and we have a single solution $\hat{\psi}_+(s_n, x) \propto \hat{\psi}_-(s_n, x) \equiv \hat{\psi}_*(s_n, s)$."

Solution for KG equation: Asymptotic solution

Asymptotic solutions ($V(x) \rightarrow 0$ as $x \rightarrow \pm\infty$)

$$[\partial_x^2 - s^2]\hat{\psi}_{\pm}(s, x) = 0 \quad (63)$$

$$\hat{\psi}_{-}(s, x) \simeq \begin{cases} e^{sx} & (x \rightarrow -\infty) \\ a_1(s)e^{sx} + a_2(s)e^{-sx} & (x \rightarrow +\infty) \end{cases}, \quad (64)$$

$$\hat{\psi}_{+}(s, x) \simeq \begin{cases} b_1(s)e^{sx} + b_2(s)e^{-sx} & (x \rightarrow -\infty) \\ e^{-sx} & (x \rightarrow +\infty) \end{cases}, \quad (65)$$

$$\mathcal{W}(s) = -2sb_2(s) \quad (x \rightarrow -\infty), \quad \mathcal{W}(s) = -2sa_1(s) \quad (x \rightarrow \infty) \quad (66)$$

$$\implies a_1(s) = b_2(s) \quad (67)$$

$$\mathcal{W}(s_n) = 0 \quad \rightarrow \quad a_1(s_n) = b_2(s_n) = 0 \quad (68)$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\langle \psi_n | \psi_m \rangle_{\text{KG}} = \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \quad (\text{Not orthogonal}) \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \quad (\text{Not orthogonal}) \\ (n = m) &= -\frac{1}{2} (\omega_n^* + \omega_n) e^{i(\omega_n^* - \omega_n)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_n x} d\Sigma^t, \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \quad (\text{Not orthogonal}) \\ (n = m) &= -\frac{1}{2} (\omega_n^* + \omega_n) e^{i(\omega_n^* - \omega_n)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_n x} d\Sigma^t, \\ (\omega_n = a_n + ib_n) &= -\frac{1}{2} 2a_n e^{2b_n t} \int_{\Sigma} e^{-2b_n x} d\Sigma^t \rightarrow \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \quad (\text{Not orthogonal}) \\ (n = m) &= -\frac{1}{2} (\omega_n^* + \omega_n) e^{i(\omega_n^* - \omega_n)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_n x} d\Sigma^t, \\ (\omega_n = a_n + ib_n) &= -\frac{1}{2} 2a_n e^{2b_n t} \int_{\Sigma} e^{-2b_n x} d\Sigma^t \rightarrow \text{diverge } (b_n < 0) \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \quad (\text{Not orthogonal}) \\ (n = m) &= -\frac{1}{2} (\omega_n^* + \omega_n) e^{i(\omega_n^* - \omega_n)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_n x} d\Sigma^t, \\ (\omega_n = a_n + ib_n) &= -\frac{1}{2} 2a_n e^{2b_n t} \int_{\Sigma} e^{-2b_n x} d\Sigma^t \rightarrow \text{diverge } (b_n < 0) \quad (\text{No E. conservation}) \end{aligned}$$

KG inner product

when $x \rightarrow \infty$,

$$\psi_n(x) = e^{-i\omega_n t} \Phi_n(x), \quad \psi_m(x) = e^{-i\omega_m t} \Phi_m(x)$$

where $s_n = -i\omega_n$, $\Phi_n(x) = e^{-sx}$

$$\begin{aligned} \langle \psi_n | \psi_m \rangle_{\text{KG}} &= \frac{i}{2} \int_{\Sigma} \left[(D_{\mu} \psi_n)^* \psi_m - \psi_n^* (D_{\mu} \psi_m) \right] d\Sigma^{\mu} \\ &= \frac{i}{2} \int_{\Sigma} \left[\left(\partial_t e^{-i\omega_n t} \Phi_n(x) \right)^* e^{-i\omega_m t} \Phi_m(x) - \left(e^{-i\omega_n t} \Phi_n(x) \right)^* \partial_t e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= \frac{i}{2} \int_{\Sigma} \left[i\omega_n^* e^{i\omega_n^* t} \Phi_n^*(x) e^{-i\omega_m t} \Phi_m(x) - e^{i\omega_n^* t} \Phi_n^*(x) (-i\omega_m) e^{-i\omega_m t} \Phi_m(x) \right] d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} \Phi_n^*(x) \Phi_m d\Sigma^t \\ &= -\frac{1}{2} (\omega_n^* + \omega_m) e^{i(\omega_n^* - \omega_m)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_m x} d\Sigma^t \neq \delta_{nm} \quad (\text{Not orthogonal}) \\ (n = m) &= -\frac{1}{2} (\omega_n^* + \omega_n) e^{i(\omega_n^* - \omega_n)t} \int_{\Sigma} e^{-i\omega_n^* x} e^{i\omega_n x} d\Sigma^t, \\ (\omega_n = a_n + ib_n) &= -\frac{1}{2} 2a_n e^{2b_n t} \int_{\Sigma} e^{-2b_n x} d\Sigma^t \rightarrow \text{diverge } (b_n < 0) \quad (\text{No E. conservation}) \end{aligned}$$

In QM,

$$\langle \psi_n | \psi_m \rangle = \delta(\omega_n - \omega_m) = \delta_{nm}$$

Solution for KG equation: Green's function method

$$\frac{d^2}{dr_*^2} \Phi''(r) + (\omega^2 - V_{\text{eff}}) \Phi(r) = 0, \quad V_{\text{eff}} = A(r) \left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} \right)$$

Solution for KG equation: Green's function method

$$\frac{d^2}{dr_*^2} \Phi''(r) + (\omega^2 - V_{\text{eff}}) \Phi(r) = 0, \quad V_{\text{eff}} = A(r) \left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} \right)$$

- This form is known as a generalized spheroidal wave equation

Solution for KG equation: Green's function method

$$\frac{d^2}{dr_*^2} \Phi''(r) + (\omega^2 - V_{\text{eff}}) \Phi(r) = 0, \quad V_{\text{eff}} = A(r) \left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} \right)$$

- ▶ This form is known as a generalized spheroidal wave equation
- ▶ their solutions are known exactly such as

$$\hat{\phi}_+(s, r) = (2is)^s e^{i\phi_+} (1 - 1/r)^s \sum_{L=-\infty}^{\infty} b_L [G_{L+\nu}(-is, isr) + iF_{L+\nu}(-is, isr)]$$

$$\hat{\phi}_-(s, r) = r^{-2s} (r - 1)^s e^{-s(r-2)} \sum_{n=0}^{\infty} a_n (1 - 1/r)^n$$

where $G_{L+\nu}$ and $F_{L+\nu}$ are Coulomb wave functions. $\phi_-(s, r)$ is analytic in s , but $\phi_+(s, r)$ has a branch cut in the complex s -plane.

Solution for KG equation: Green's function method


$$\frac{d^2}{dr_*^2} \Phi''(r) + (\omega^2 - V_{\text{eff}}) \Phi(r) = 0, \quad V_{\text{eff}} = A(r) \left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} \right)$$

- ▶ This form is known as a generalized spheroidal wave equation
- ▶ their solutions are known exactly such as

$$\hat{\phi}_+(s, r) = (2is)^s e^{i\phi_+} (1 - 1/r)^s \sum_{L=-\infty}^{\infty} b_L [G_{L+\nu}(-is, isr) + iF_{L+\nu}(-is, isr)]$$

$$\hat{\phi}_-(s, r) = r^{-2s} (r - 1)^s e^{-s(r-2)} \sum_{n=0}^{\infty} a_n (1 - 1/r)^n$$

where $G_{L+\nu}$ and $F_{L+\nu}$ are Coulomb wave functions. $\phi_-(s, r)$ is analytic in s , but $\phi_+(s, r)$ has a branch cut in the complex s -plane.



Solution for KG equation: Green's function method


$$\frac{d^2}{dr_*^2} \Phi''(r) + (\omega^2 - V_{\text{eff}}) \Phi(r) = 0, \quad V_{\text{eff}} = A(r) \left(\frac{l(l+1)}{r^2} + \frac{A'(r)}{r} \right)$$

- ▶ This form is known as a generalized spheroidal wave equation
- ▶ their solutions are known exactly such as

$$\hat{\phi}_+(s, r) = (2is)^s e^{i\phi} + (1 - 1/r)^s \sum_{L=-\infty}^{\infty} b_L [G_{L+\nu}(-is, isr) + iF_{L+\nu}(-is, isr)]$$

$$\hat{\phi}_-(s, r) = r^{-2s} (r - 1)^s e^{-s(r-2)} \sum_{n=0}^{\infty} a_n (1 - 1/r)^n$$


where $G_{L+\nu}$ and $F_{L+\nu}$ are Coulomb wave functions. $\phi_-(s, r)$ is analytic in s , but $\phi_+(s, r)$ has a branch cut in the complex s -plane.



Solution for KG equation: Green's function method

$$\hat{\psi}(s, x) = \int_{-\infty}^{\infty} dx' \hat{G}(s, x, x') \mathcal{J}(s, x), \quad (69)$$

$$G(t, x, x') = \int_{\epsilon - i\infty}^{\epsilon + i\infty} \frac{ds}{2\pi i} e^{st} \hat{G}(s, x, x') \quad (70)$$



$$G(t, x, x') = G_{\text{QNM}}(t, x, x') + G_{\text{B}}(t, x, x') + G_{\text{F}}(t, x, x') \quad (71)$$

Branch cut solution for small ω : late time tail

- In the limit $t \rightarrow \infty$ with x fixed :

$$G_B(t, x, x') \simeq (-1)^{l+1} \frac{2(2l+2)!}{[(2l+1)!!]^2} \frac{r_h}{c} \frac{(xx')^{l+1}}{t^{2l+3}} \quad (72)$$

there is a power-law tail at spatial infinity, i.e. a non-radiative tail.


- In the limit $t \rightarrow \infty$ with t/x fixed :

$$G_B(t, x, x') \simeq (-1)^{l+1} \frac{(l+1)!}{(2l+1)!!} \frac{r_h}{c} \frac{(x')^{l+1}}{u^{l+2}}, \quad u \equiv t - x \quad (73)$$

there is a radiative tail at future null infinity.

Solution for KG equation: Green's function method

$$G(t, x, x') = G_{\text{QNM}}(t, x, x') + G_{\text{B}}(t, x, x') + G_{\text{F}}(t, x, x') \quad (74)$$



- ▶ $t \simeq x - x'$: the outward propagating wave can effectively ignore the influence of this potential and travel outward reaches the observer, G_{F}
- ▶ $t \simeq x + x'$: (quasinormal modes) the part of the low-frequency scalar perturbation traveling towards the black hole bounces off the effective potential $V_{\text{eff}}(r)$ near the horizon and propagates back to the observer, G_{QNM}
- ▶ $t \gg x + x'$: (late-time behavior) the QNMs gradually fade away, and only the decaying modes of the signal persist. The outgoing wave that traveled to a large distance and then scattered back towards the observer due to the spacetime curvature at large distances, G_{B}

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation
- ▶ Hermitian system
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation vs energy is not conserved (open system)
- ▶ observable is real (e.g. $E \sim \omega$ is real)
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation vs energy is not conserved (open system)
- ▶ observable is real (e.g. $E \sim \omega$ is real) vs ω is complex
- ▶ eigenfunctions are orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation vs energy is not conserved (open system)
- ▶ observable is real (e.g. $E \sim \omega$ is real) vs ω is complex
- ▶ eigenfunctions are orthonormal vs eigenfunctions are not orthonormal
- ▶ eigenfunctions form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation vs energy is not conserved (open system)
- ▶ observable is real (e.g. $E \sim \omega$ is real) vs ω is complex
- ▶ eigenfunctions are orthonormal vs eigenfunctions are not orthonormal
- ▶ eigenfunctions form a complete set vs eigenfunctions do not form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable)

Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation vs energy is not conserved (open system)
- ▶ observable is real (e.g. $E \sim \omega$ is real) vs ω is complex
- ▶ eigenfunctions are orthonormal vs eigenfunctions are not orthonormal
- ▶ eigenfunctions form a complete set vs eigenfunctions do not form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable) vs no Hilbert space (KG inner product is not well defined and not square-integrable)


Summary

In QM (vs in QNMs),

- ▶ solve Schrodinger equation vs solve KG equation (but, Schrodinger equation like)
- ▶ Hermitian system vs non-Hermitian
- ▶ energy conservation vs energy is not conserved (open system)
- ▶ observable is real (e.g. $E \sim \omega$ is real) vs ω is complex
- ▶ eigenfunctions are orthonormal vs eigenfunctions are not orthonormal
- ▶ eigenfunctions form a complete set vs eigenfunctions do not form a complete set
- ▶ Hilbert space (inner product is well defined and square-integrable) vs no Hilbert space (KG inner product is not well defined and not square-integrable)
- ▶ Despite QNMs not being a Hermitian system themselves, numerous developments have been made to utilize the powerful mathematical properties of Hermitian systems.

Black Hole ringdwon

Ring-down phase



The newly formed, perturbed black hole relaxes to a stable state (usually a Kerr black hole characterized by its mass M and spin J) by emitting gravitational waves. These waves are described by the black hole's quasinormal modes (QNMs)

QNMs for gravitational wave

We need to consider metric perturbation

$$g_{\alpha\beta}(x) = \bar{g}_{\alpha\beta}(x) + h_{\alpha\beta}(x) \quad (75)$$

- ▶ for a Schwarzschild black hole
 - ▶ Odd-parity (axial) modes: Use Regge-Wheeler equation, related to off-diagonal metric components.
 - ▶ Even-parity (polar) modes: Use Zerilli equation, related to diagonal metric components.
- ▶ for a Kerr black hole : Teukolsky equation

They all take a form of

$$\frac{d^2}{dr_*^2} \Phi''(r) + (\omega^2 - V) \Phi(r) = 0 \quad (76)$$

Ringdown phase with QNMs for gravitational wave

However, as we have seen that *QNMs generally do not form a complete set in the usual mathematical sense due to their non-Hermitian nature* and the boundary conditions imposed. They can be part of a generalized basis when combined with other modes, such as:

$$\psi(t,x) = \sum_n c_n \psi_n(x) e^{-i\omega_n t} + \text{branch cut contribution} \quad (77)$$

²Kostas D. Kokkotas, Bernd G. Schmidt, "Quasi-Normal Modes of Stars and Black Holes", Volume 2, article number 2, (1999)

Ringdown phase with QNMs for gravitational wave

However, as we have seen that *QNMs generally do not form a complete set in the usual mathematical sense due to their non-Hermitian nature* and the boundary conditions imposed. They can be part of a generalized basis when combined with other modes, such as:

$$\psi(t,x) = \sum_n c_n \psi_n(x) e^{-i\omega_n t} + \text{branch cut contribution} \quad (77)$$




Figure 4: The response of a Schwarzschild black hole as a Gaussian wave packet impinges upon it. The QNM signal dominates the signal after $t \approx 70M$ while at later times (after $t \approx 300M$) the signal is dominated by a power-law fall-off with time.

Thank you!